



# Delaware

## Live births, abortions, and pregnancies

### Live birth rates, 1982–1992



### Live birth rates,\* 1991



### Pregnancy rates, 1990

*Pregnancy rates are not available because Delaware did not report abortion data to CDC.*

### National live birth rate ranking, 1990


Age group		
15-17	18-19	15-19
20th highest	37th highest	29th highest

\*Missing bar indicates fewer than 10,000 females or fewer than 20 births. This graph refers to females aged 15–19 years.




# Sexually transmitted diseases Delaware

## Gonorrhea rates, 1982–1992



## Gonorrhea rates, 1992



## National gonorrhea rate ranking,\* 1992


Males – 12th highest    Females – 10th highest

\*National ranking refers to males and females aged 10–24 years.




## Chlamydia cases and rates,\* 1988–1992


Cases



Rates




Cases



Rates



## Acquired immunodeficiency syndrome (AIDS) cumulative cases,† 1981–1992



\*Chlamydia cases and rates refer to males and females of all ages.


†Missing bar indicates fewer than six cumulative cases.




# Related risk behaviors,\* 1993

Delaware


Had sexual intercourse (lifetime)




Had sexual intercourse with four or more people (lifetime)



Had sexual intercourse during the past three months (currently sexually active)




Used a condom during last sexual intercourse (among currently sexually active)



Used birth control pills during last sexual intercourse (among currently sexually active)



Injected illegal drugs (lifetime)



\*Related risk behaviors refer to high school students.

Sources: CDC—NCHS (birth data), NCPS (gonorrhea and chlamydia data), NCID (AIDS data); 1993 Delaware Youth Risk Behavior Survey (risk behavior data).

